CONTENTS

(Abstracts/contents lists published in: Chem. Abstr.; Curr. Contents, Phys. Chem. Earth Sci., Life Sci.)

Die Kristall- und Molekülstrukturen von drei thermodynamisch benachteiligten 2,6-Anhydro-nitroalditolen ("Glycopyranosylnitromethanen")
J. Kopf, M. Morf, B. Zimmer, O. Jarchow (Hamburg, B.R.D.), H. Brandenburg, und P. Köll (Oldenburg, B.R.D.) 1

A reverse approach to ^{1}H-n.m.r. assignments of bacterial polysaccharides
D. N. M. Jones and J. K. M. Sanders (Cambridge, Gt. Britain) 15

(1-^{13}C) Alditols: elimination of magnetic equivalence in ^{1}H- and ^{13}C-n.m.r. spectra of symmetric compounds through (^{13}C)-substitution
E. C. Garrett and A. S. Serianni (Notre Dame, IN, U.S.A.) 23

Synthesis of a tetrasaccharide of the genus-specific lipopolysaccharide epitope of *Chlamydia*
P. Kosma, R. Bahnmüller, G. Schulz (Vienna, Austria), and H. Brade (Borstel, F.R.G.) . 37

Synthesis of the sodium salts of methyl 2-O-α-L-fucopyranosyl-α-L-fucopyranoside 3- and 4-sulfate
R. K. Jain and K. L. Matta (Buffalo, NY, U.S.A.) 51

Synthesis of disaccharide methyl glycosides related to the polysaccharide from *Klebsiella* serotype 40 and a study of their inhibition in the preciptin reaction
U. B. Maddali, A. K. Ray, and N. Roy (Calcutta, India) 59

Synthesis of 4-*O*-methyl-β-rhodomycins using derivatives of 4-amino-4-deoxy- and 3,4-diamino-3,4-dideoxy sugars
C. Kolar, K. Dehmel, and H. Moldenhauer (Marburg, F.R.G.) 67

Synthesis and conformational studies of carrabiose and its 4′-sulphate and 2,4′-disulphate
E. Parra, H.-N. Caro, J. Jiménez-Barbero, M. Martín-Lomas, and M. Bernabé (Madrid, Spain) 83

Polysaccharides from thermal polymerization of glucosides
G. R. Ponder and G. N. Richards (Missoula, MT, U.S.A.) 93

Preparative separation of *myo*-inositol bis- and tris-phosphate isomers by anion-exchange chromatography
B. Goldschimdt (Perstorp, Sweden) 105

Anthracycline oligosaccharides: facile stereoselective synthesis of 2,6-dideoxy-α-L-*lyxo*-hexopyranosides
C. Kolar, G. Kneissl, H. Wolf, and T. Kämpchen (Marburg, F.R.G.) 111

ROESY studies and HSEA calculations on xylose-containing oligosaccharides related to *N*-glycoproteins
B. R. Leeflang, J. B. Bouwstra, J. Kerékgyártó, J. P. Kamerling, and J. F. G. Vliegenthart (Utrecht, The Netherlands) 117

$^{87}Rb^{+}$ spin relaxation in enzymically purified and in untreated iota-carrageenan
L. Piculell (Lund, Sweden) and C. Rochas (Grenoble, France) 127

Structure of the K24 antigen of *E. coli* 083:K24:H, a polymer that consists of α-Kdo*p* and glycerol phosphate
M. C. Lenter, B. Jann, and K. Jann (Freiburg-Zähringen, F.R.G.) 139

Structural studies of the capsular polysaccharide from *Aerococcus viridans* var. *homari*
K. Hermansson, L. Kenne, B. Lindberg (Stockholm, Sweden), B. Arie, R. G. Brown, and J. E. Stewart (Halifax, N.S., Canada) 145

Structure of a unique sulfated α-L-galactofucan from the tunicate *Clavelina*
M. S. G. Pavão, P. A. S. Mourão (Rio de Janeiro, Brazil), and B. Mulloy (Potters Bar, Gt. Britain) 153

Structural studies of a sulfated L-galactan from *Styela plicata* (Tunicate): analysis of the Smith-degraded polysaccharide
R. M. Albano, M. S. G. Pavão, P. A. S. Mourão (Rio de Janeiro, Brazil), and B. Mulloy (Potters Bar, Gt. Britain) 163

Oxidation of oligoglycosyl alditols during methylation catalyzed by sodium hydroxide and iodomethane in methyl sulfoxide
W. S. York, L. L. Kiefer, P. Albersheim, and A. G. Darvill (Athens, GA, U.S.A.) 175

Competitive formation of cellulose *p*-toluenesulfonate and chlorodeoxycellulose during homogeneous reaction of *p*-toluenesulfonyl chloride with cellulose in *N*,*N*-dimethylacetamide–lithium chloride
C. L. McCormick, T. R. Dawsey, and J. K. Newman (Hattiesburg, MS, U.S.A.) 183

The structural analysis of a levan produced by *Streptococcus salivarius* SS2
P. J. Simms, W. J. Boyko, and J. R. Edwards (Villanova, PA, U.S.A.) 193

Solute effects on the thermal stability of glycerol monostearate–amylose complex superstructures
C. G. Biliaderis and H. D. Seneviratne (Winnipeg, Man., Canada) 199

Molecular structure of 3-*O*-(3,6-anhydro-α-D-galactopyranosyl)-β-D-galactopyranose (neocarrabiose) in the solid state and in solution: an investigation by X-ray crystallography, n.m.r. spectroscopy, and molecular mechanics calculations
D. Lamba, A. L. Segre (Rome, Italy), S. Glover, W. Mackie, B. Sheldrick (Leeds, Gt. Britain), and S. Pérez (Nantes, France) 215

Notes

X-Ray crystallographic and n.m.r. (^{1}H and ^{13}C) studies of methyl 4,6-*O*-(*R*)-benzylidene-3-*O*-methyl-2-*O*-*p*-tolylsulfonyl-α-D-glucopyranoside and methyl 4,6-*O*-(*R*)-benzylidene-2-*O*-methyl-3-*O*-*p*-tolylsulfonyl-β-D-glucopyranoside
E. Lee, P. McArdle, N. Melody, D. Cunningham, and J. Gallagher (Galway, Ireland) . . 231

Synthesis of methyl 2-*O*-α-D-galactopyranosyl-3-*O*-β-D-glucopyranosyl-α-D-glucopyranoside, a trisaccharide of the R-1 core antigen of *Enterobacteriaceae*
S. Basu and J. N. Pal (Calcutta, India) 241

3-Deoxy-L-*glycero*-pentos-2-ulose (3-deoxy-L-xylosone) and L-*threo*-pentos-2-ulose (L-xylosone) as intermediates in the degradation of L-ascorbic acid
D. B. Shin and M. S. Feather (Columbia, MO, U.S.A.) 246

Phase-transfer-catalyzed synthesis of a phenylseleno α-ketoside of *N*-acetylneuraminic acid, (phenyl 5-acetamido-3,5-dideoxy-D-*glycero*-α-D-*galacto*-2-selenononulopyranosid)onic acid; a new sialidase inhibitor
J. Rothermel and H. Faillard (Saarbrücken, F.R.G.) 251

Chiral recognition of 1-*O*-allyl- and 1-*O*-benzyl-D- and -L-*myo*-inositol by cyclomalto-hexaose and -heptaose (α- and β-cyclodextrin)
M. Bernabé, J. Jiménez-Barbero, M. Martin-Lomas, S. Penadés, and C. Vicent (Madrid, Spain) 255

Efficient syntheses of 3,4,6-tri-*O*-acetyl-2-deoxy-2-phthalimido-β- and -α-D-galactopyranosyl chloride
U. Nilsson, A. K. Ray, and G. Magnusson (Lund, Sweden) 260

A simple and ready internal *C*-glycosylation of 2,3,5-tri-*O*-benzylglycofuranoses promoted by the boron trifluoride–diethyl ether complex
M. Anastasia, P. Allevi, P. Ciuffreda, A. Fiecchi, and A. Scala (Milan, Italy) 264

Synthesis and immunoadjuvant activity of *N*-[2-*O*-(2-acetamido-1,2,3,5-tetradeoxy-1,5-imino-D-glucitol-3-yl)-D-lactoyl]-L-alanyl-D-isoglutamine
H. Ishida, M. Kitagawa, M. Kiso, A. Hasegawa (Gifu, Japan), and I. Azuma (Sapporo, Japan) 267

Some studies concerning the dimerization of 2,3-*O*-isopropylidene-L-erythrose
L. M. Lerner (Brooklyn, NY, U.S.A.) 273

3-Deoxy-D-*erythro*-hexulose: a convenient synthesis and its interaction with the enzymes of fructose metabolism
W. L. Dills, Jr. (North Dartmouth, MA, U.S.A.) 276

Synthesis of methyl 2-*O*-(sodium α-L-fucopyranosyl 3- and 4-sulfate)-α-L-fucopyranoside
R. K. Jain and K. L. Matta (Buffalo, NY, U.S.A.) 280

A one-step synthesis of 6-*S*-acetyl-6-deoxy-1,2:3,4-di-*O*-isopropylidene-6-thio-α-D-galactopyranose
M. von Itzstein, M. J. Jenkins, and M. Mocerino (Parkville, Vic., Australia) 287

Colorimetric determination of alginates and fragments thereof as the 2-nitrophenylhydrazides
Y. Murata, S. Kawashima, E. Miyamoto, N. Masauji, and A. Honda (Kanozawa, Japan) 289

Structural studies of the *O*-specific polysaccharide from *Salmonella kentucky* strain 98/39 (O:8,H:*i*,Z_6)
V. I. Torgov, V. N. Shibaev, A. A. Shashkov, and S. Sh. Rozhnova (Moscow, U.S.S.R.) . 293

Structure of 3-*O*-(α-D-glucopyranosyluronic acid)-L-galactopyranose, an aldobiouronic acid isolated from the polysaccharides of various unicellular red algae
S. Geresh, O. Dubinsky, S. Arad, D. Christiaen, and R. Glaser (Beersheva, Israel) 301

The effects of deglycosylation on the properties of native and biotinylated bovine testicular hyaluronidase
D. Lace, A. H. Olavesen, and P. Gacesa (Cardiff, Gt. Britain) 306

Modified malto-oligosaccharides as inhibitors of human alpha-amylases
K. Omichi, S. Hase, and T. Ikenaka (Osaka, Japan) 312

Errata c1

Author index c3

Subject index c5